



Assessment of Aleutian mink disease virus (AMDV) prevalence in feral American mink in Iceland. Case study of a pending epizootiological concern in Europe

Remigiusz Panicz¹, Piotr Eljasik¹, Jakub Skorupski², Przemysław Śmietana², Róbert A. Stefánsson³, Menja von Schmalensee³ and Magdalena Szenejko²

¹Department of Meat Technology, Faculty of Food Sciences and Fisheries, West Pomeranian University of Technology Szczecin, Szczecin, Poland

²Institute of Marine and Environmental Sciences, Molecular Biology and Biotechnology Centre, University of Szczecin, Szczecin, Poland

³West Iceland Nature Research Centre, Stykkisholmur, Iceland

ABSTRACT

Background. Recurring escapes or deliberate releases and subsequent infiltration or establishment of feral populations by individuals from fur farms have been commonly noted since the beginning of fur industry expansion. Once animals have invaded ecosystems adjacent to source farms escapees can change the demography of the feral populations through hybridization, outbreeding depression, competition and spreading of various pathogens which can decimate wild populations. In our study, we aimed to assess spread of Aleutian mink disease virus (AMDV) in the feral population of American mink (*Neovison vison*) in Iceland. The additional objective was to elucidate whether basic morpho-anatomical parameters (*i.e.*, Fulton's condition factor or spleen to body weight ratio) might be used as a preliminary indicator of AMDV infection.

Methods. American mink ($n = 164$) were captured by professional hunters in 8 regions of Iceland. The detection of AMDV in the spleen of male and female individuals was based on PCR amplification of an NS1 gene fragment.

Results. We confirmed AMDV presence in 23.8% ($n = 39$) of collected samples with no significant difference in infection rate between males and females. Additionally, we revealed that the prevalence of virus in the feral population was higher closer to fur farms. However, the countrywide prevalence and direction of AMDV distribution needs to be further investigated. Comparison of condition indices in non-infected and infected animals showed significant deterioration of body and spleen parameters in the latter group. Therefore, the application of basic measurements of the American mink may be used to evaluate the health status of individuals in terms of pathogen infection.

Conclusions. The study shed a new light on prevalence and distribution of AMDV in the feral population of American mink in Iceland and the results might be successfully applied to develop models to infer dynamics of various pathogens, even those latently transmitted by disease-free animals.

Submitted 22 March 2021
Accepted 4 August 2021
Published 17 September 2021

Corresponding author
Remigiusz Panicz, rpanicz@zut.edu.pl

Academic editor
Xavier Harrison

Additional Information and
Declarations can be found on
page 13

DOI 10.7717/peerj.12060

© Copyright
2021 Panicz et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Ecology, Molecular Biology, Veterinary Medicine, Virology, Zoology

Keywords American mink, Feral population, Free-ranging population, Fulton's condition factor, Spleen, Virus prevalence and spread

INTRODUCTION

The American mink (*Neovison vison*) is the main species farmed by the fur industry in Iceland since farms were first established in 1931 ([Hersteinsson, 1992](#); [Hersteinsson, 1999](#); [Bonesi & Palazon, 2007](#)). In the following years, excluding the period 1951–1969 when fur-farming was banned by law, events of mink escapes have been recorded ([Skírnisson, Gunnarsson & Hjartardóttir, 1990](#)). A countrywide feral population had been established by 1975 ([Stefansson, Schmalensee & Skorupski, 2016](#)). Escape events have been confirmed also in e.g., Norway ([Bevanger & Henriksen, 1995](#)), Denmark ([Pertoldi et al., 2013](#)), Scotland ([Macdonald et al., 2015](#)) and France ([Fournier-Chambrillon et al., 2004](#)) where American mink, after accidental escapes or deliberate releases, quickly established viable, feral and invasive populations. In the early years of the mink farming industry in Iceland, mink escapes were observed and probably quite common, but the frequency of such events in recent times should be evaluated in the further studies ([Stefansson, Schmalensee & Skorupski, 2016](#)). The success of animals in the wild, evidenced by rapid increase of feral populations, has been attributed to the species' high reproduction capability, opportunistic foraging and the frequent lack of natural enemies and competitors in invaded areas ([Bevanger & Henriksen, 1995](#); [Maran et al., 2016](#)). Spread of non-indigenous American mink is associated with a number of serious ecological threats, such as intensive foraging (e.g., small mammals, like *Arvicola amphibious*, ground-nesting birds, fish), competition and aggression (e.g., *Mustela lutreola*) against indigenous fauna ([Heggenes & Borgstrom, 1988](#); [Macdonald & Strachan, 1999](#); [Macdonald & Harrington, 2003](#); [Nordström et al., 2003](#); [Banks et al., 2008](#)). Interestingly, the escapee events may threaten native wild American mink populations in their natural range, through hybridization, outbreeding depression, and competition for food, space, and mates ([Bowman et al., 2007](#)).

Farmed American mink are usually housed in relatively high densities, which favours multiplication and spread of pathogens. Therefore, domestic mink which escape from farms can transmit Aleutian mink disease virus (AMDV) and other pathogens (e.g., zoonotic *Cryptosporidium* sp.) that may seriously diminish reproduction and survival rate of feral mink populations ([Nituch et al., 2011](#); [Qian et al., 2020](#)). *N. vison* is also reported to be susceptible to coronaviruses ([Vlasova et al., 2011](#)). Recently, American mink were identified as hosts of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), the causative agent of deadly for humans COVID-19 disease ([Hossain, Akter & Saha, 2020](#)). The transmission of SARS-CoV-2 to American mink was confirmed in several countries (e.g., in the Netherlands and Denmark), where increased mortality of animals was observed ([Molenaar et al., 2020](#); [Maron, 2020](#)).

Among several disease conditions described for American mink, currently the most significant one is Aleutian mink disease (AMD), a highly pathogenic parvovirus affecting American mink and other mustelids ([Bloom, Kanno & Mori, 1994](#); [Virtanen et al., 2020](#)).

The AMD transmission is horizontal (*via* blood, saliva, faeces or urine) or vertical (seropositive dams during the perinatal period infects kits) (Alexandersen *et al.*, 1985) and clinical symptoms in adult mink include hypergammaglobulinemia, kidneys glomeruli inflammation, reduced fertility, and spontaneous abortion (Alexandersen, 1986; Bloom, Kanno & Mori, 1994). Occasionally AMD leads to neurologic disease associated with a nonsuppurative meningoencephalitis (Jahns *et al.*, 2010). Kits affected by AMD show rapidly progressing interstitial pneumonia, extensive atelectasis and hyaline membrane formation, which results in high mortality rates. Depending on strain, mortality rate varies from 30% to over 90%, and adult survivors develop typical lesions of adult form (Alexandersen, 1986).

AMD is caused by Aleutian mink disease virus, a single-stranded DNA parvovirus of the *Amdoparvovirus* genus (Cotmore *et al.*, 2014). Similarly to other parvoviruses, AMDV replicates using rolling hairpin mechanism (Cotmore & Tattersall, 1996). However, AMDV can be distinguished from other parvoviruses by a host response to exposure, which contributes to the progression of disease symptoms (Castelruiz, Blixenkroner-Møller & Aasted, 2005). The virus has a genome of approx. 4.8 kbp and non-enveloped virion contains two structural (VP1 and VP2) and three non-structural (NS1, NS2 and NS3) proteins (Bloom *et al.*, 1988). Among these proteins NS1 and VP2 are suggested to have crucial roles in pathogenicity, since a high degree of variability in both amino acid sequences between highly virulent and non-virulent strains was recorded (Best, Wolfenbarger & Bloom, 2002).

AMDV has frequently been reported in feral and wild American mink populations in Europe and North America. Yamaguchi & Macdonald (2001) found that 52% of investigated feral individuals (only adults) in the upper Thames region of the UK were anti-AMDV antibodies positive. In France, 23% of caught free-ranging American mink were infected with AMDV (Fournier-Chambrillon *et al.*, 2004), while in a Spanish feral population 40% of tested individuals of *N. vison* were infected (Mañas *et al.*, 2001). Wild populations were also tested in *N. vison*'s native range, *i.e.*, Nova Scotia (Canada), where 93% of collected samples were positive (Farid, 2013).

Isolated islands are interesting areas for virus studies. Jensen *et al.* (2012) showed high, up to 45%, occurrence of AMDV in feral American mink in Bornholm (Denmark). Additionally, virus strains were found to be clustered into two groups, suggesting introduction of Swedish and Danish strains to Bornholm (Jensen *et al.*, 2012). In Iceland, samples from 1986–87 showed minimal (3.5%) prevalence of AMDV in the feral population (Skírnisson, Gunnarsson & Hjartardóttir, 1990). Moreover, Ryt-Hansen *et al.* (2017) proved that 4 sequences of partial NS1 gene from Icelandic free-ranging mink were close to the Swedish virus strain.

Iceland, due to its insulated character, provides a unique opportunity to accurately monitor spread and persistence of pathogens in feral populations. Therefore, the overarching objective of this study was to assess the prevalence of AMDV in the feral population of American mink in Iceland. Additionally, we aimed to address the question whether AMDV-infected American mink in the wild exhibit deterioration in their health status.

MATERIALS & METHODS

Sample collection, morphometric analysis and tissue sampling

Feral mink were hunted by professional mink hunters using hunting dogs or death traps as part of an official control program funded by Icelandic authorities. A total of 164 (100 males and 64 females) American mink carcasses of mink caught in 2010–2018 in 8 regions were obtained for post-mortem analyses and sample collection performed at the West Iceland Nature Research Centre. Body weight to the nearest 1 g and total body length (excluding tail) to the nearest 0.5 cm were assessed and sex was determined. The entire spleen was removed and weighed to the nearest 0.01 g. Spleen samples were packed individually in sterile bags and stored at -20°C until delivery to the Faculty of Food Science and Fisheries of the West Pomeranian University of Technology (Szczecin) molecular laboratory for later DNA extraction and analysis.

AMDV detection

Spleen samples were used for DNA extraction which was performed using the High Pure PCR Template Preparation Kit (Roche) according to the manufacturer's instructions. The quantity and quality of DNA isolates were assessed using a NanoDrop 2000 (Thermo Scientific) and by separation in 1.5% agarose gel. The extracted DNA was tested using end-point PCR and unique set of primers to amplify 374 bp fragment of the AMDV NS1 gene: AMDV-F-7-H-PN1: 5'-CATATTCAGTGTGCTTAGGTTA-3' and AMDV-R-7-HPN2: 5'-CGTTCCTTTGTTAGTTAGGTTGTC-3' (Jensen *et al.*, 2011). Amplifications were conducted on Mastercycler (Eppendorf) under following conditions: 1 step of 5 min at 94°C followed by 45 cycles at 94°C for 30 s, 55°C for 30 s, 72°C for 30 s, and final extension at 72°C for 7 min. Reactions were prepared with the GoTaq PCR kit (Promega), including 5 μl of Green GoTaq[®] Flexi Buffer, 2.5 μl of MgCl_2 (25 mM Solution), 0.5 μl of PCR Nucleotide Mix (10 mM), 0.125 μl of GoTaq[®] DNA Polymerase (5 u/l), 0.5 μM of each primer and 3 μl of DNA template in final volume of 25 μl . Negative sample was applied to ensure specificity of PCR reaction. The results of end-point PCR reactions were assessed by 2.0% gel electrophoresis and positive samples sequenced on both strands by means of direct Sanger sequencing by the Genomed company (Warsaw) to eliminate false-positive results. Raw sequences were processed, aligned and checked using the sequence editing software Geneious v.11.1.5 (created by Biomatters, available from <http://www.geneious.com>). Additionally, all sequences were translated into amino acids to confirm their accuracy and to detect the presence of nuclear DNA pseudogenes, insertions, deletions or stop codons. Finally, all sequences were searched for matches against the GeneBank database using BLASTn and number of haplotypes were calculated using the DNAsp 6.0 software (Rozas *et al.*, 2017).

Condition index, statistical analysis and data visualization

Clear sexual dimorphism of examined morphometric parameters was observed. Therefore, Fulton's condition factor (K) and the ratio of spleen weight to body weight were used for the comparison of the potential impact of viral infection. Fulton's condition factor was calculated with the following formula: $K = 100 \times W \times L_b^{-3}$, where K is Fulton's




Figure 1 The eight sampling regions (light red) in Iceland. Sample sites are indicated with red diamonds (AMDV positive) and blue dots (AMDV negative). Mink farms in operation in 2012 are shown in black. The number of AMDV positive samples/total number of samples are indicated for each region.

Full-size [DOI: 10.7717/peerj.12060/fig-1](https://doi.org/10.7717/peerj.12060/fig-1)

condition factor, W is body weight (g) and L_b is body length (cm). A Mann–Whitney U test was used to check significance of differences between Fulton’s condition indexes, spleen weight and body weight of infected and AMDV-free males and females and pairwise comparisons between groups were conducted to correct for multiple testing (post-hoc tests). All sampled individuals from across Iceland as was treated as exchangeable. The theoretical linear distance between sampling site and nearest mink farm was measured based on [Stefansson, Schmalensee & Skorupski \(2016\)](#) data ([Fig. 1](#)). That distance is likely to differ from a true distance travelled by feral mink in the area, since American mink migrate mainly along waterways and coastline ([Macdonald et al., 2015](#)). The simplification was implemented assuming the same systematic error for all the samples, which allows to correctly interpret the results. The Median test ([Zar, 2010](#)) was used to compare distance from sampling site to the nearest mink farm between AMDV-infected and virus-free individuals. The map was produced using the QGIS 3.12 software (QGIS Development Team, available from <http://qgis.osgeo.org>). Statistical analyses were performed using the Statistica software (Statsoft) and data were visualised using R version 4.0.3 ([R Core Team, 2020](#)) and R-package ggplot2 ([Wickham, 2016](#)).




Figure 2 (A) Distribution of AMDV positive individuals ($n = 39$) collected in eight regions of Iceland in sex groups. (B) Comparison of AMDV infection rate (diameter of the circles) in both sexes. An analysis of Pearson's Chi-squared test with Yates' continuity correction ($\chi^2 = 1.9559$, $df = 1$, p -value = 0.162) indicated non-significant difference.

Full-size DOI: 10.7717/peerj.12060/fig-2

RESULTS

Virus detection

Molecular analysis showed that 23.8% of collected samples were AMDV positive (39 out of 164) and 33 haplotypes were identified using the DNAsp 6.0 software ([Supplementary Information S1](#)). AMDV haplotypes sequencing data are available at GenBank under accession numbers [MW772385–MW772417](#). While most of the animals sampled in our study were males ([Fig. 2A](#)), Pearson's χ^2 test with Yates' continuity correction revealed no significant difference in infection rate between males and females ([Fig. 2B](#)). Statistical analysis of distances of sample locations relative to mink farm location ([Stefansson, Schmalensee & Skorupski, 2016](#)) revealed that AMDV infected American mink were on average closer (40.8 km) to farms than disease free mink (117.4 km), ([Figs. 1 and 3](#)).

Morphometry and condition index

Results of statistical analysis of studied morphometric features revealed significantly higher body weight (t -test; $t = 11.653$, $df = 145$; $p < 0.0001$) and body length (t -test; $t = 15.186$; $df = 133$; $p < 0.001$) of males comparing to females. Moreover, study showed low values of the variation coefficient (less than 10%) for the mean body length of both males and females ([Table 1](#)). Comparison of Fulton's condition factor (K) between non-infected and infected individuals revealed significantly higher value of this indicator in non-infected males and females. Moreover, infected males had significantly lower K than infected females and now difference in K was found between AMDV-free males and females ([Fig. 4](#)). Additionally, the study revealed significantly higher spleen weight to body weight ratio in males and females infected with AMDV, and higher ratio in females comparing to males, regardless of the infection ([Fig. 5](#)).




Figure 3 Comparison of distance between nearest farm and AMDV infected or non-infected American mink individuals. The Pearson's Chi-squared test ($\chi^2 = 29.388$, $df = 1$, $p = 0.0000$) indicates a significant difference.

Full-size DOI: 10.7717/peerj.12060/fig-3

DISCUSSION

Virus detection

Isolated islands are unique places for virological studies, especially because of the limited influx of virus strains from the outside. Here, we confirmed presence of AMDV in free-ranging American mink in 6 (Austur-Húnavatnssýsla, Borgarfjar arsýsla, Dalasýsla, Nor ur-Múlasýsla, Su ur-Múlasýsla and Vestur-Húnavatnssýsla) out of 8 investigated regions (Fig. 1). AMDV was not detected in two regions (Nor ur-Ísafjar arsýsla, Snæfellsnes - og Hnappadalssýsla). However, to confirm our outcomes further research should focus on different disease tracking methods in such areas. For instance, non-invasive methods such as collecting faeces combined with a sensitive detection method, *i.e.*, qPCR (Virtanen *et al.*, 2020). This approach has been successfully used for the detection of amdoparvoviruses in farmed American mink (Virtanen *et al.*, 2020), raccoon dog (*Nyctereutes procyonoides*) and Arctic fox (*Vulpes lagopus*) (Shao *et al.*, 2014) or wild red fox (*Vulpes vulpes*) (Bodewes *et al.*, 2014).

The study showed that 23.8% of collected mink samples were found AMDV positive and the result is consistent with a previous study of Ryt-Hansen *et al.* (2017), who reported AMDV in 25% (4 out of 16) of samples collected in Iceland, however, no information on sampling sites were provided. These results are similar to estimates in France where AMDV prevalence amounted to 23% (Fournier-Chambrillon *et al.*, 2004), but lower than in other

Table 1 Basic morphometric parameters taken from all *N. vison* individuals.

Sex	Mean		Standard deviation		Lower 95% conf. limit		Higher 95% conf. limit		Coef. of variation %	
	Male	Female	Male	Female	Male	Female	Male	Female	Male	Female
Body length (excluding tail) (cm)	41.2	35.2	2.46	1.54	40.6	34.72	41.7	35.60	5.98	4.38
Body weight (g)	1173.3	728.9	258.99	136.53	1121.1	692.72	1225.5	765.17	22.07	18.73
Spleen weight (g)	2.9381	2.3922	1.4838	1.1086	2.6358	2.1058	3.2404	2.6785	50.50	46.34
Fulton's condition factor <i>K</i>	1.67	1.67	0.3132	0.339	1.61	1.58	1.74	1.77	18.73	20.24
Spleen weight/body weight ratio	0.0026	0.0033	0.0012	0.0015	0.002	0.0029	0.003	0.0038	46.79	45.55




Figure 4 Evaluation of Fulton's condition factor (K) for mink in Iceland using Mann-Whitney U test. Comparison of Fulton's condition factor between specimens of mink: infected and non-infected males ($Z = 3.183$); infected and non-infected females ($Z = 3.183$); non-infected males and non-infected females ($Z = -0.159$); infected males and infected females ($Z = 2.576$).

Full-size DOI: 10.7717/peerj.12060/fig-4

countries where the feral American mink is considered an invasive species, *i.e.*, Sweden (Persson *et al.*, 2015), Denmark (Jensen *et al.*, 2012), and Spain (Mañas *et al.*, 2001), where AMDV prevalence amounted to 57%, 45% and 40% respectively. The dominance of male individuals (in collected and AMDV-positive samples) may be related to hunting methods used in this study, since they have been found to be more susceptible to trapping, because of wider home-range size (Buskirk & Lindstedt, 1989). Thus, considering similar infection rate of males and females, the main type of AMDV transmission in Icelandic free-ranging American mink remains unclear. According to Alexandersen *et al.* (1985) and Macdonald *et al.* (2015) horizontal transmission occurs mainly in males, due to roam-and-mate tactics, but vertical transmission from dams to kits during the perinatal period were equally plausible in both sexes. Additionally, our results contrast with the hypothesis that in mammals male individuals are more susceptible to pathogens, since revealed infection rate was similar for both males and females (Moore & Wilson, 2002).

The results indicated that AMDV infected American mink were on average closer to farms than disease free individuals. Nituch *et al.* (2011) reported similar findings with a higher prevalence of infected mink near farms in a large-scale study conducted in Canada, which indicated mink farms as a source of AMDV spread in the wild population




Figure 5 Evaluation of spleen weight/body weight ratio for mink in Iceland using Mann–Whitney U test. Comparisons of spleen weight/ body weight ratio between specimens of mink: infected and non-infected males ($Z = -2.935$), infected and non-infected females ($Z = -2.586$), non-infected males and non-infected females ($Z = -3.915$), infected males and infected females ($Z = 2.403$).

Full-size [DOI: 10.7717/peerj.12060/fig-5](https://doi.org/10.7717/peerj.12060/fig-5)

of American mink. In some countries, such a distribution might be explained by newly escaped, infected farmed mink tending to occupy areas close to the farm they escaped from (Ionescu *et al.*, 2018). Such an explanation is however highly unlikely in this study. Icelandic feral mink are distinctively different from farmed mink in body size and fur quality (Stefansson, Schmalensee & Skorupski, 2016), and farm escapees are thus very easily recognized. During the sampling from mink carcasses used in this study, every animal was assessed as a part of the feral population (pers. comm. Stefansson and von Schmalensee). It should be noted that in general, feral or wild American mink may also transmit pathogens to domestic mink, and at least one such scenario has been reported (Nituch *et al.*, 2012). But the occurrence of pathogen transmission between free-ranging and domestic mink needs to be investigated further. The frequency of spill-back events may eventually depend on many various local factors, such as the infection rate in the free-ranging mink and their population density near mink farms, the biosecurity measures taken at farms but also natural behaviour during the mating season, since they display scramble competition polygyny *i.e.*, competitive mate search (Alcock, 1998).

This study sheds light on the prevalence of AMDV in parts of Iceland but does not provide conclusive evidence about the direction of AMDV transmission, as samples were

not obtained throughout whole country, nor from mink farms. Further studies should address possible factors which may influence the main routes of AMDV transmission, especially historical background, specific for Iceland. When AMDV entered the feral mink population in this country in 1970–1983, the seropositive farms were all located in the northern part of the island. The highest proportion of seropositive feral mink in 1986–87 were also located in N-Iceland (*Skírnisson, Gunnarsson & Hjartardóttir, 1990*), and the highest infection rate in feral mink has continued to be connected to N-Iceland (*Stefánsson et al., 2004; Stefánsson et al., 2011*).

From the ecological point of view, the biggest concern in Iceland is the potential risk of AMDV transmission to Arctic fox (*Vulpes lagopus*), the only native carnivore in Iceland (*Unnsteinsdóttir et al., 2016*). AMDV has been detected in red foxes (*Vulpes vulpes*), proving that AMDV is capable of infecting canids, but to assess the virus threat for *V. lagopus* further investigations are needed (*Canuti et al., 2020a*).

In light of the COVID-19 pandemic and reported circulation of SARS-CoV-2 virus between human and American mink (*Maron, 2020*) our results may indicate how virus can spillover to the natural environment in isolated islands, such as Iceland. However, from the early views the transmission of SARS-CoV-2 appears to be much more rapid (*Molenaar et al., 2020*) than AMDV, which is transmitted poorly and can be controlled in farm conditions (*James & Dubovi, 2016*). Therefore, sealing the American mink farms and minimization of escapees is critical to both stop viruses spread and avoid the need for radical moves, like culling of the entire populations in Netherlands and Denmark (*Molenaar et al., 2020; Maron, 2020*).

Morphometry and condition index

The low values of the variation coefficient (less than 10%) for the mean body length of both males and females are an interesting aspect of the results. The high variability of mean values of other analysed parameters indicates an existence of a sex independent separating factor. Comparisons showed significantly higher body weight and body length of males comparing to females that is consistent with the sexual dimorphism in the species (e.g., *Kruska & Schreiber, 1999*). Considering that the separating factor, mentioned above, does not differentiate individuals with respect to body length, but other morphometric indices studied, suggests that it is a pathogenic factor, possibly AMDV. This factor divides the studied population into at least two statistical subpopulations, within both males and females.

The observed sexual dimorphism is typical for this species (at all development stages) and has been previously reported elsewhere (*Melero et al., 2012; Macdonald et al., 2015*). Moreover, male and female American minks' body length coefficient of variation indicates a homogeneous population (in statistical terms) relative to this feature. Assuming that body length is a relatively stable feature in adult organisms, high coefficient of variation values of other features indicates influence of differentiating factors. In further analysis we considered AMDV infection as such factor since its impact on *N. vison* health status and condition was previously reported (*Eklund et al., 1968*).

Condition indices are a simple method to investigate influence of external factors on an organism, and therefore its health status (Stevenson & Woods Jr, 2006). Fulton's condition factor (K) is generally used in fishery studies, but the potential of its application in small mammals has been reported (Peig & Green, 2010). In the study non-infected mink had significantly higher value of K comparing to AMDV-infected individuals. Jensen, Chriél & Hansen (2015) showed worsening condition of American mink infected with AMDV in the experimental challenges. Our results showed similar tendency in free-ranging and adjacent to farms population of American mink, regardless of sex. The worse condition of infected *N. vison* individuals could be explained with symptoms (anaemia, anorexia) described for AMD (Eklund et al., 1968). In contrast, Canuti et al. (2020b) revealed no association between AMDV infection and body condition in Pacific marten (*Martes caurina*) and the North American river otter (*Lontra canadensis*), although it is unclear whether these species develop disease symptoms or act as asymptomatic carriers. Additionally, study found that only infected males had significantly lower K than infected females. Thus, indicating a stronger negative impact on males compared to females, despite similar infection rate. To our knowledge, no reports on the sexual disparities in response to AMDV infection have been published elsewhere, however during an American mink population decrease in Iceland (2004–2010) male individuals were significantly more impaired than females (Magnusdottir et al., 2014). An analogous pattern was reported in house mouse (*Mus musculus*) infected with influenza virus (Celestino et al., 2018), while male performed better than female in the South American coati (*Nasua nasua*) infected with trypanosomatids and gastrointestinal parasites (Olifiers et al., 2015).

The size of immune organs (*i.e.*, spleen, performing important immune functions) has been commonly used to describe animals' investment in immunological response to various factors (John, 1994; Schulte-Hostedde, Bowman & Nituch, 2012). In the study AMDV infected mink had significantly higher spleen weight to body weight ratio. The enlargement of spleen was most likely caused by pathogen presence, which means immune organs were in a highly activated (primed) state and increased number of lymphocytes (*e.g.*, T-cells) were produced (Abolins et al., 2017). The spleen enlargement of AMDV infected mink is consistent with previous findings (Reichert & Kostro, 2014). Our results were also comparable to findings of Persson et al. (2015), where difference in relative spleen weight was found in male free-ranging American mink, but no difference was found in females, probably due to a smaller sample size. Additionally, regardless of the infection females had higher spleen weight to body weight ratio than males. The difference of relative spleen weight between sexes has been reported elsewhere (Schulte-Hostedde, Bowman & Nituch, 2012; Persson et al., 2015), but the reason for sexual disparity remains unclear. Moore & Wilson (2002) suggested a shift of energy resources towards reproduction and/or growth over the immune system in male mammals and immunosuppressive side effects of androgenic hormones. While Nunn et al. (2009) found no differences in immune system functioning associated with sexual dimorphism. Therefore, males' lower value of relative spleen weight reported in our study should be further investigated. Moreover, we propose Fulton's condition factor combined with spleen weight to body weight ratio to be

a preliminary measure to conduct further testing for AMDV infection in American mink using combination of more sensitive methods.

CONCLUSIONS

The results of our study provide a simple method to preliminarily assess the possibility of an AMDV infection based on simple and quick body measurements of American mink. The reported distribution and prevalence of AMDV may be used as starting point for further studies on prevalence and transmission of different pathogens (e.g., SARS-CoV-2, *Cryptosporidium* sp.) in Iceland and other isolated ecosystems. Of particular importance is to study feral and wild American mink as potential reservoirs of SARS-CoV-2, as possibility of infection to American mink has been proved (Molenaar et al., 2020; Maron, 2020). Further studies should focus on assessment of AMDV prevalence in the remaining counties in Iceland and identification of other pathogens that might be transmitted in a latent form by healthy animals. In general, monitoring of diseases in free-ranging populations in the vicinity of mink farms should be commenced world-wide, to evaluate both the prevalence of pathogen spill-over from farms and the risk of pathogen spill-back from feral or wild populations.

ACKNOWLEDGEMENTS

This work was supported by the West Iceland Nature Research Centre, Stykkisholmur, Iceland; they coordinated and performed sampling, including shipment of samples to Poland. This study would not have been possible without the effort made by Icelandic mink hunters, who provided us with samples of their catch.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

The work was supported by the Financial Mechanism of the European Economic Area (EEA) 2009–2014 under Grant number 15/2017/Wn16/OP-XN-02/D and the National Fund for Environmental Protection and Water Management (Poland) under Grant number 17/2017/Wn16/OP-XN-02/D. The Department of Meat Sciences (Faculty of Food Science and Fisheries, West Pomeranian University of Technology, Szczecin) and the Institute of Marine and Environmental Sciences, and Molecular Biology and Biotechnology Centre, Szczecin, Poland (University of Szczecin) provided all necessary resources (access to the laboratory, financial support) to perform molecular, bioinformatic and graphical tasks. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

Financial Mechanism of the European Economic Area (EEA) 2009–2014: 15/2017/Wn16/OP-XN-02/D.

National Fund for Environmental Protection and Water Management (Poland):
17/2017/Wn16/OP-XN-02/D.

Faculty of Food Science and Fisheries, West Pomeranian University of Technology,
Szczecin.

Institute of Marine and Environmental Sciences.

Molecular Biology and Biotechnology Centre, Szczecin, Poland.

Competing Interests

The authors declare there are no competing interests.

Author Contributions

- Remigiusz Panicz, Jakub Skorupski and Róbert A. Stefánsson conceived and designed the experiments, performed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Piotr Eljasik and Przemysław Śmietana conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.
- Menja von Schmalensee conceived and designed the experiments, performed the experiments, authored or reviewed drafts of the paper, and approved the final draft.
- Magdalena Szenejko analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.

Data Availability

The following information was supplied regarding data availability:

The raw data are available in the [Supplementary File](#). The sequences are available in the Supplemental File and at GenBank: [MW772385–MW772417](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.12060#supplemental-information>.

REFERENCES

- Abolins S, King EC, Lazarou L, Weldon L, Hughes L, Drescher P, Raynes JG, Hafalla JCR, Viney ME, Riley EM. 2017.** The comparative immunology of wild and laboratory mice, *Mus musculus domesticus*. *Nature Communications* **8**:1–13
[DOI 10.1038/ncomms14811](https://doi.org/10.1038/ncomms14811).
- Alcock J. 1998.** *Animal behaviour: an evolutionary approach*. 10th edition. Sunderland: Sinauer Associates, Inc.
- Alexandersen S. 1986.** Acute interstitial pneumonia in mink kits: experimental reproduction of the disease. *Veterinary Pathology* **23**:579–588
[DOI 10.1177/030098588602300506](https://doi.org/10.1177/030098588602300506).
- Alexandersen S, Uttenthal-Jensen A, Hansen M, Aasted B. 1985.** Experimental transmission of Aleutian disease virus (ADV) to different animal species. *Acta*

Pathologica Microbiologica ScandInavica Series B: Microbiology **93**:195–200
DOI [10.1111/j.1699-0463.1985.tb02876](https://doi.org/10.1111/j.1699-0463.1985.tb02876).

- Banks PB, Nordstrom M, Ahola M, Salo P, Fey K, Korpimaki E. 2008.** Impacts of alien mink predation on island vertebrate communities of the Baltic Sea Archipelago: review of a long-term experimental study. *Boreal Environment Research* **13**:3–16.
- Best SM, Wolfenbarger JB, Bloom ME. 2002.** Caspase activation is required for permissive replication of Aleutian mink disease parvovirus in vitro. *Virology* **292**:224–234
DOI [10.1006/viro.2001.1238](https://doi.org/10.1006/viro.2001.1238).
- Bevanger K, Henriksen G. 1995.** The distributional history and present status of the American mink (*Neovison vison*, Schreber, 1777) in Norway. *Annales Zoologici Fennici* **32**:11–14
DOI [10.18276/ab.2016.23-08](https://doi.org/10.18276/ab.2016.23-08).
- Bloom ME, Kaaden OR, Huggans E, Cohn A, Wolfenbarger JB. 1988.** Molecular comparisons of in vivo- and in vitro-derived strains of Aleutian disease of mink parvovirus. *Journal of Virology* **62**:132–138
DOI [10.1128/JVI.62.1.132-138.1988](https://doi.org/10.1128/JVI.62.1.132-138.1988).
- Bloom ME, Kanno H, Mori S. 1994.** Aleutian mink disease: puzzles and paradigms. *Infectious Agents and Disease* **3**:279–301.
- Bodewes R, Ruiz-Gonzalez A, Schapendonk CM, vanBrand JM, Osterhaus AD, Smits SL. 2014.** Viral metagenomic analysis of feces of wild small carnivores. *Virology Journal* **11**:89
DOI [10.1186/1743-422X-11-89](https://doi.org/10.1186/1743-422X-11-89).
- Bonesi L, Palazon S. 2007.** The American mink in Europe: status, impacts, and control. *Biological Conservation* **134**:470–483
DOI [10.1016/j.biocon.2006.09.006](https://doi.org/10.1016/j.biocon.2006.09.006).
- Bowman J, Kidd AG, Gorman RM, Schulte-Hostedde AI. 2007.** Assessing the potential for impacts by feral mink on wild mink in Canada. *Biological Conservation* **139**:12–18
DOI [10.1016/j.biocon.2007.05.020](https://doi.org/10.1016/j.biocon.2007.05.020).
- Buskirk SW, Lindstedt SL. 1989.** Sex biases in trapped samples of Mustelidae. *Journal of Mammalogy* **70**:88–97
DOI [10.2307/1381672](https://doi.org/10.2307/1381672).
- Canuti M, McDonald E, Graham SM, Rodrigues B, Bouchard É, Neville R, Pitcher M, Whitney HG, Marshall HD, Lang AS. 2020a.** Multi-host dispersal of known and novel carnivore amdoxparvoviruses. *Virus Evolution* **veaa072**
DOI [10.1093/ve/veaa072](https://doi.org/10.1093/ve/veaa072).
- Canuti M, Todd M, Monteiro P, Van Osch K, Weir R, Schwantje H, Britton AP, Lang AS. 2020b.** Ecology and infection dynamics of multi-host amdoxparvoviral and protoparvoviral carnivore pathogens. *Pathogens* **9**(62):124
DOI [10.3390/pathogens9020124](https://doi.org/10.3390/pathogens9020124).
- Castelruiz Y, Blixenkroner-Moller M, Aasted B. 2005.** DNA vaccination with the Aleutian mink disease virus NS1 gene confers partial protection against disease. *Vaccine* **23**:1225–1231
DOI [10.1016/j.vaccine.2004.09.003](https://doi.org/10.1016/j.vaccine.2004.09.003).
- Celestino I, Checconi P, Amatore D, De Angelis M, Coluccio P, Dattilo R, Fegatelli DA, Clemente AM, Matarrese P, Torcia MG, Mancinelli R, Mammola CL, Garaci E, Vestri AR, Malorni W, Palamara AT, Mancinelli R. 2018.** Differential redox state contributes to sex disparities in the response to influenza virus infection in male and female mice. *Frontiers in Immunology* **9**:1747
DOI [10.3389/fimmu.2018.01747](https://doi.org/10.3389/fimmu.2018.01747).

- Cotmore SF, Agbandje-McKenna M, Chiorini JA, Mukha DV, Pintel DJ, Qiu J, Soderlund-Venermo M, Tattersall P, Tijssen P, Gatherer D, Davison AJ. 2014. The family parvoviridae. *Archives of Virology* **159**:1239–1247
DOI [10.1007/s00705-013-1914-1](https://doi.org/10.1007/s00705-013-1914-1).
- Cotmore SF, Tattersall P. 1996. Parvovirus DNA replication. In: DePhamphilis ML, ed. *DNA replication in eukaryotic cells*. New York: Cold Spring Harbor Laboratory Press, 799–813.
- Eklund CM, Hadlow WJ, Kennedy RC, Boyle CC, Jackson TA. 1968. Aleutian Disease of Mink: Properties of the Etiologic Agent and the Host Responses. *The Journal of Infectious Diseases* **118**(5):510–526.
- Farid AH. 2013. Aleutian mink disease virus in furbearing mammals in Nova Scotia, Canada. *Acta Veterinaria Scandinavica* **55**:1–9 DOI [10.1186/1751-0147-55-10](https://doi.org/10.1186/1751-0147-55-10).
- Fournier-Chambrillon C, Aasted B, Perrot A, Pontier D, Sauvage F, Artois M, Cassiède J-M, Chauby X, Molin AD, Simon C, Fournier P. 2004. Antibodies to Aleutian mink disease parvovirus in free-ranging European mink (*Mustela lutreola*) and other small carnivores from southwestern France. *Journal of Wildlife Diseases* **40**:394–402
DOI [10.7589/0090-3558-40.3.394](https://doi.org/10.7589/0090-3558-40.3.394).
- Heggenes J, Borgstrom R. 1988. Effect of mink, *Mustela vison* Schreber, predation on cohorts of juvenile Atlantic salmon, *Salmo salar* L. and brown trout, *Salmo trutta* L. in 3 small streams. *Journal of Fish Biology* **33**:885–894
DOI [10.1111/j.1095-8649.1988.tb05536.x](https://doi.org/10.1111/j.1095-8649.1988.tb05536.x).
- Hersteinsson P. 1992. Mammals of the Thingvallavatn area. *Oikos* **64**:396–404
DOI [10.2307/3545061](https://doi.org/10.2307/3545061).
- Hersteinsson P. 1999. Methods to eradicate the American mink (*Mustela vison*) in Iceland. Workshop on the control and eradication of non-native terrestrial vertebrates. *Report to the Council of Europe T-PVS (99)* **26**:17–21.
- Hossain MG, Akter S, Saha S. 2020. SARS-CoV-2 host diversity: an update of natural infections and experimental evidences. *Journal of Microbiology, Immunology and Infection* DOI [10.1016/j.jmii.2020.06.006](https://doi.org/10.1016/j.jmii.2020.06.006).
- Ionescu DT, Hodor C, Druga M, Druga M. 2018. Recent occurrence of the American mink (*Neovison vison*) in the central Romania. In: *Proceedings of the biennial international symposium forest and sustainable development*. Brasov, Romania:.
- Jahns H, Daly P, McElroy MC, Sammin DJ, Bassett HF, Callanan JJ. 2010. Neuropathologic features of Aleutian disease in farmed mink in Ireland and molecular characterization of Aleutian mink disease virus detected in brain tissues. *Journal of Veterinary Diagnostic Investigation* **22**:101–105 DOI [10.1177/104063871002200120](https://doi.org/10.1177/104063871002200120).
- James NM, Dubovi EJ. 2016. *Veterinary virology*. Cambridge: Academic Press
DOI [10.1016/B978-0-12-800946-8.00007-6](https://doi.org/10.1016/B978-0-12-800946-8.00007-6).
- Jensen TH, Chriél M, Hansen MS. 2015. Progression of experimental chronic Aleutian mink disease virus infection. *Acta Veterinaria Scandinavica* **58**:35
DOI [10.1186/s13028-016-0214-7](https://doi.org/10.1186/s13028-016-0214-7).
- Jensen TH, Christensen LS, Chriél M, Uttenthal Å, Hammer AS. 2011. Implementation and validation of a sensitive PCR detection method in the eradication campaign

- against Aleutian mink disease virus. *Journal of Virological Methods* 171:81–85 DOI 10.1016/j.jviromet.2010.10.004.
- Jensen TH, Christensen LS, Chriél M, Harslund J, Salomonsen CM, Hammer AS. 2012.** High prevalence of Aleutian mink disease virus in free-ranging mink on a remote Danish island. *Journal of Wildlife Diseases* 48:497–502 DOI 10.7589/0090-3558-48.2.497.
- John JL. 1994.** The avian spleen: a neglected organ. *Quarterly Review of Biology* 69:327–351 DOI 10.1086/418649.
- Kruska D, Schreiber A. 1999.** Comparative morphometrical and biochemical-genetic investigations in wild and ranch mink (*Mustela vison*: Carnivora: Mammalia). *Acta Theriologica* 44(4):377–392.
- Mañas S, Ceña JC, Ruiz-Olmo J, Palazón S, Domingo M, Wolfenbarger JB, Bloom ME. 2001.** Aleutian mink disease parvovirus in wild riparian carnivores in Spain. *Journal of Wildlife Diseases* 37:138–144 DOI 10.7589/0090-3558-37.1.138.
- Macdonald DW, Harrington LA. 2003.** The American mink: the triumph and tragedy of adaptation out of context. *New Zealand Journal of Zoology* 30:421–441 DOI 10.1080/03014223.2003.9518350.
- Macdonald DW, Harrington LA, Yamaguchi N, Thom MDF, Bagniewska J. 2015.** Biology, ecology, and reproduction of American mink *Neovison vison* on lowland farmland. In: Macdonald DW, Feber RE, eds. *Wildlife conservation on farmland: conflict in the countryside*. Vol. 12. Oxford: Oxford University Press, 127–147.
- Macdonald DW, Strachan R. 1999.** *The mink and the water vole: analyses for conservation*. Oxford: WildCRU.
- Magnusdottir R, Schmalensee Mvon, Stefansson RA, Macdonald DW, Hersteinsson P. 2014.** A foe in woe: American mink (*Neovison vison*) diet changes during a population decrease. *Mammalian Biology* 79:58–63 DOI 10.1016/j.mambio.2013.08.002.
- Maran T, Skumatov D, Gomez A, dra MPo, Abramov AV, Dinets V. 2016.** *Mustela lutreola* *Mustela lutreola*. The IUCN Red List of Threatened Species 2016: e.T14018A45199861.http://dx.doi.org/10.2305/IUCN.UK.2016-1.RLTS.T14018A45199861.en. (accessed on 22 November 2020).
- Maron DF. 2020.** Denmark to cull 15 million mink after coronavirus spillover into humans. National Geographic. Available at <https://www.nationalgeographic.com/animals/2020/11/denmark-mink-culling/> (accessed on 08 November 2020)).
- Melero Y, Santulli G, Gómez A, Gosálbez J, Rodriguez-Refojos C, Palazón S. 2012.** Morphological variation of introduced species: the case of American mink (*Neovison vison*) in Spain. *Mammalian Biology* 77:345–350 DOI 10.1016/j.mambio.2012.02.001.
- Molenaar RJ, Vreman S, Hakze-van der Honing RW, Zwart R, De Rond J, Weesendorp E, Smit LAM, Koopmans M, Bouwstra R, Stegeman A, Vander Poel WH. 2020.** Clinical and pathological findings in SARS-CoV-2 disease outbreaks in farmed mink (*Neovison vison*). *Veterinary Pathology* 57:653–657 DOI 10.1177/0300985820943535.
- Moore SL, Wilson K. 2002.** Parasites as a viability cost of sexual selection in natural populations of mammals. *Science* 297(5589):2015–2018 DOI 10.1126/science.1074196.

- Nituch LA, Bowman J, Beauclerc KB, Schulte-Hostedde AI. 2011. Mink farms predict Aleutian disease exposure in wild American mink. *PLOS ONE* 6(67):e21693 DOI 10.1371/journal.pone.0021693.
- Nituch LA, Bowman J, Wilson P, Schulte-Hostedde AI. 2012. Molecular epidemiology of Aleutian disease virus in free-ranging domestic, hybrid, and wild mink. *Evolutionary Applications* 5:330–340 DOI 10.1111/j.1752-4571.2011.00224.x.
- Nordström M, Högmänder J, Jarmo L, Nummelin J, Laanetu N, Korpimäki E. 2003. Effects of feral mink removal on seabirds, waders and passerines on small islands in the Baltic Sea. *Biological Conservation* 109:359–368 DOI 10.1016/S0006-3207(02)00162-3.
- Nunn CL, Lindenfors P, Pursall ER, Rolff J. 2009. On sexual dimorphism in immune function. *Philosophical Transactions of the Royal Society B. Biological Sciences* 364(1513):61–69 DOI 10.1098/rstb.2008.0148.
- Olifiers N, Jansen AM, Herrera HM, Bianchi RDC, D’Andrea PS, Mourão GDM, Gompper ME. 2015. Co-infection and wild animal health: effects of trypanosomatids and gastrointestinal parasites on coatis of the Brazilian Pantanal. *PLOS ONE* 10(612):e0143997 DOI 10.1371/journal.pone.0143997.
- Peig J, Green AJ. 2010. The paradigm of body condition: a critical reappraisal of current methods based on mass and length. *Functional Ecology* 24:1323–1332 DOI 10.1111/j.1365-2435.2010.01751.x.
- Persson S, Jensen TH, Blomström AL, Appelberg MT, Magnusson U. 2015. Aleutian mink disease virus in free-ranging mink from Sweden. *PLOS ONE* 10(63):e0122194 DOI 10.1371/journal.pone.0122194.
- Pertoldi C, Rodjajn S, Zalewski A, Demontis D, Loeschke V, Kjaersgaard A. 2013. Population viability analysis of American mink (*Neovison vison*) escaped from Danish mink farms. *Journal of Animal Science* 91:2530–2541 DOI 10.2527/jas.2012-6039.
- Qian W, Zhang Y, Jiang Y, Zhao A, Lv C, Qi M. 2020. Molecular characterization of *Cryptosporidium* spp. in minks (*Neovison vison*), blue foxes (*Vulpes lagopus*), and raccoon dogs (*Nyctereutes procyonoides*) in farms from Xinjiang, Northwest China. *Parasitology Research* 119(611):3923–3927 DOI 10.1007/s00436-020-06909-8.
- R Core Team. 2020. *R: a language and environment for statistical computing*. Vienna: R Foundation for Statistical Computing. Available at <https://www.R-project.org/>.
- Reichert M, Kostro K. 2014. Effect of persistent infection of mink with Aleutian mink disease virus on reproductive failure. *Bulletin of the Veterinary Institute in Pulawy* 58:369–373 DOI 10.2478/bvip-2014-0057.
- Rozas J, Ferrer-Mat A, Sánchez-De I, Barrio JC, Guirao-Rico S, Librado P, Ramos-Onsins SE, Sánchez-Gracia A. 2017. DnaSP 6: DNA sequence polymorphism analysis of large datasets. *Molecular Biology and Evolution* 34:3299–3302 DOI 10.1093/molbev/msx248.
- Ryt-Hansen P, Hagberg EE, Chriél M, Struve T, Pedersen AG, Larsen LE, Hjulsgager CK. 2017. Global phylogenetic analysis of contemporary aleutian mink disease viruses (AMDVs). *Virology Journal* 14:1–6 DOI 10.1186/s12985-017-0898-y.

- Schulte-Hostedde AI, Bowman J, Nituch LA. 2012.** Dynamic spleen mass in wild and domestic American mink. *Biological Journal of the Linnean Society* **107**:624–631 DOI [10.1111/j.1095-8312.2012.01961.x](https://doi.org/10.1111/j.1095-8312.2012.01961.x).
- Shao XQ, Wen YJ, Ba HX, Zhang XT, Yue ZG, Wang KJ, Li CY, Qiu J, Yang FH. 2014.** Novel amdoparvovirus infecting farmed raccoon dogs and arctic foxes. *Emerging Infectious Diseases* **20**:2085 DOI [10.3201/eid2012.140289](https://doi.org/10.3201/eid2012.140289).
- Skírnisson K, Gunnarsson E, Hjartardóttir S. 1990.** Plasmacytosis-sýking í villtum mink á Íslandi. *Icelandic Agricultural Science* **3**:113–122.
- Stefánsson RA, Bjarnadóttir S, Hersteinsson P, Gunnarsson E. 2004.** Útbreiddsla veirusjúkdómsins plasmacytosis Á villta minkastofninum [Distribution of Aleutian mink disease virus in the feral mink population in Iceland]. 2011. Abstract of a poster presented at the Science symposium, Reykjavik 16-17 April 2004. Abstract book, p. 116. [In Icelandic].
- Stefánsson RA, Von Schmalensee M, Gunnarsson E, Hersteinsson P. 2011.** Veirusjúkdómurinn plasmacytosis í villtum mink. [Aleutian mink disease virus in feral mink in Iceland]. Abstract of a presentation at the Icelandic Biological Conference, Reykjavik 11-12 November 2011. [In Icelandic].
- Stefánsson RA, Schmalensee M, von, Skorupski J. 2016.** A tale of conquest and crisis: invasion history and status of the American mink (*Neovison vison*) in Iceland. *Acta Biologica* **23**:87–100 DOI [10.18276/ab.2016.23-08](https://doi.org/10.18276/ab.2016.23-08).
- Stevenson RD, Woods Jr WA. 2006.** Condition indices for conservation: new uses for evolving tools. *Integrative & Comparative Biology* **46**:1169–1190 DOI [10.1093/icb/icl052](https://doi.org/10.1093/icb/icl052).
- Unnsteinsdóttir ER, Hersteinsson P, Pálsson S, Angerbjörn A. 2016.** The fall and rise of the Icelandic Arctic fox (*Vulpes lagopus*): a 50-year demographic study on a noncyclic Arctic fox population. *Oecologia* **181**:1129–1138 DOI [10.1007/s00442-016-3635-0](https://doi.org/10.1007/s00442-016-3635-0).
- Virtanen J, Aaltonen K, Vapalahti O, Sironen T. 2020.** Development and validation of nucleic acid tests to diagnose Aleutian mink disease virus. *Journal of Virology Methods* **279**:113776 DOI [10.1016/j.jviromet.2019.113776](https://doi.org/10.1016/j.jviromet.2019.113776).
- Vlasova AN, Halpin R, Wang S, Ghedin E, Spiro DJ, Saif LJ. 2011.** Molecular characterization of a new species in the genus Alphacoronavirus associated with mink epizootic catarrhal gastroenteritis. *Journal of General Virology* **92(66)**:1369 DOI [10.1099/vir.0.025353-0](https://doi.org/10.1099/vir.0.025353-0).
- Wickham H. 2016.** *ggplot2: elegant graphics for data analysis*. New York: Springer-Verlag. Available at <https://ggplot2.tidyverse.org>.
- Yamaguchi N, Macdonald DW. 2001.** Detection of Aleutian disease antibodies in feral American mink in southern England. *Veterinary Record* **149**:485–488 DOI [10.1136/vr.149.16.485](https://doi.org/10.1136/vr.149.16.485).
- Zar JH. 2010.** *Biostatistical analysis*. 5th edition. Illinois: Northern Illinois University Pearson.